



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Macrophage 11-HSD1 deficiency promotes inflammatory angiogenesis

Citation for published version:

Zhang, Z, Coutinho, AE, Man, JTY, Kipari, TMJ, Hadoke, P, Salter, DM, Seckl, JR & Chapman, K 2017, 'Macrophage 11-HSD1 deficiency promotes inflammatory angiogenesis', *Journal of Endocrinology*.
<https://doi.org/10.1530/JOE-17-0223>

Digital Object Identifier (DOI):

[10.1530/JOE-17-0223](https://doi.org/10.1530/JOE-17-0223)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Journal of Endocrinology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Macrophage 11 β -HSD-1 deficiency promotes inflammatory angiogenesis

Zhenguang Zhang^{1,*}, Agnes E Coutinho^{1,†}, Tak Yung Man¹, Tiina M J Kipari², Patrick W F Hadoke¹, Donald M Salter², Jonathan R Seckl¹ and Karen E Chapman¹

¹University/BHF Centre for Cardiovascular Science, The Queen's Medical Research Institute, The University of Edinburgh, Edinburgh, UK

²Centre for Genomic and Experimental Medicine, MRC Institute of Genetic and Molecular Medicine, The University of Edinburgh, Western General Hospital, Edinburgh, UK

* (Z Zhang is now at AV Hill Building, Faculty of Life Science, University of Manchester, Manchester, UK)

† (A E Coutinho is now at Humber College Institute of Technology and Advanced Learning, Toronto, Canada)

Correspondence should be addressed to K E Chapman
Email
Karen.Chapman@ed.ac.uk

Abstract

11 β -Hydroxysteroid dehydrogenase-1 (11 β -HSD1) predominantly converts inert glucocorticoids into active forms, thereby contributing to intracellular glucocorticoid levels. 11 β -HSD1 is dynamically regulated during inflammation, including in macrophages where it regulates phagocytic capacity. The resolution of inflammation in some disease models including inflammatory arthritis is impaired by 11 β -HSD1 deficiency or inhibition. However, 11 β -HSD1 deficiency/inhibition also promotes angiogenesis, which is beneficial in mouse models of surgical wound healing, myocardial infarction or obesity. The cell types responsible for the anti-inflammatory and anti-angiogenic roles of 11 β -HSD1 have not been characterised. Here, we generated *Hsd11b1*^{CKO} mice with *LysM-Cre* mediated deletion of *Hsd11b1* to investigate whether 11 β -HSD1 deficiency in myeloid phagocytes is pro-angiogenic and/or affects the resolution of inflammation. Resolution of inflammatory K/BxN-induced arthritis was impaired in *Hsd11b1*^{CKO} mice to a similar extent as in mice globally deficient in 11 β -HSD1. This was associated with >2-fold elevation in levels of the endothelial marker *Cdh5* mRNA, suggesting increased angiogenesis in joints of *Hsd11b1*^{CKO} mice following arthritis. A pro-angiogenic phenotype was confirmed by measuring angiogenesis in subcutaneously implanted polyurethane sponges, in which *Hsd11b1*^{CKO} mice showed 20% greater vessel density than their littermate controls, associated with higher expression of *Cdh5*. Thus, 11 β -HSD1 deficiency in myeloid phagocytes promotes angiogenesis. Targeting 11 β -HSD1 in macrophages may be beneficial in tissue repair.

Key Words

- ▶ steroid metabolism
- ▶ glucocorticoid
- ▶ inflammation
- ▶ 11 β -HSD1
- ▶ macrophage

Journal of Endocrinology
(2017) **234**, 291–299

Introduction

Glucocorticoids exert anti-inflammatory effects both *in vivo* and *in vitro*. Whilst the potent effects of synthetic glucocorticoids have been widely exploited clinically to treat inflammatory disease, including rheumatoid arthritis, the role of endogenous corticosteroids in regulating inflammation is less well understood (Coutinho & Chapman 2011).

Nevertheless, macrophages have emerged as a key target for the anti-inflammatory and immunomodulatory effects of endogenous glucocorticoids (Bhattacharyya *et al.* 2007, Tuckermann *et al.* 2007). Macrophages orchestrate much of the tissue remodelling that follows injury (Minutti *et al.* 2016) and play a key role in the angiogenesis that is

important for tissue repair (Nucera *et al.* 2011). Macrophages are also important in the excessive or dysregulated angiogenesis that contributes to the pathogenesis of many chronic inflammatory diseases, a maladaptive response in rheumatoid arthritis (Jackson *et al.* 1997, Koch 1998).

Macrophages express 11 β -hydroxysteroid dehydrogenase type 1 (11 β -HSD1), a glucocorticoid metabolising enzyme that, in intact cells, converts the intrinsically inert cortisone and 11-dehydrocorticosterone into the active glucocorticoids, cortisol and corticosterone, respectively (Chapman *et al.* 2013b). 11 β -HSD1 thus amplifies glucocorticoid action. 11 β -HSD1 deficiency or inhibition alters inflammatory responses (Chapman *et al.* 2013a). 11 β -HSD1-deficient mice show more severe acute inflammation in thioglycollate-induced sterile peritonitis (Gilmour *et al.* 2006), following lipopolysaccharide (Zhang & Daynes 2007), in carrageenan-induced pleurisy (Coutinho *et al.* 2012), in the K/BxN serum-induced model of inflammatory arthritis (Coutinho *et al.* 2012) and following coronary artery ligation in a model of myocardial infarction (McSweeney *et al.* 2010). Macrophages are implicated in the altered inflammatory responses of 11 β -HSD1-deficient mice. *In vitro*, peritoneal or splenic macrophages from 11 β -HSD1-deficient mice overproduce inflammatory cytokines following lipopolysaccharide stimulation (Zhang & Daynes 2007). During thioglycollate-induced peritonitis, 11 β -HSD1-deficient mice show a delay in the acquisition of phagocytic capacity by macrophages, though the inflammation resolves at a similar time to that in control mice (Gilmour *et al.* 2006). Reduced cholesterol accumulation in macrophages plays a role in the atheroprotective and anti-inflammatory effects of global 11 β -HSD1 deficiency or inhibition in atherosclerosis-prone *ApoE*^{-/-} mice (Garcia *et al.* 2013, Kipari *et al.* 2013, Luo *et al.* 2013). Following myocardial infarction, macrophages accumulate around the infarct zone more rapidly in 11 β -HSD1-deficient mice (McSweeney *et al.* 2010). They switch to a pro-reparative, pro-angiogenic 'M2' phenotype earlier than in control mice, resulting in greater angiogenesis in the healing infarct and better recovery of heart function post-myocardial infarction (Small *et al.* 2005, McSweeney *et al.* 2010, Michailidou *et al.* 2012). Global 11 β -HSD1 deficiency is also pro-angiogenic in other contexts: in the sponge implantation assay (Small *et al.* 2005); in adipose tissue of obese mice where it is associated with reduced adipose tissue hypoxia and inflammation (Michailidou *et al.* 2012) and in cutaneous surgical wounds, where it is associated with improved wound repair (Small *et al.* 2005).

Here, we have generated mice deficient in 11 β -HSD1 in myeloid phagocytes (*Hsd11b1*^{MDKO}) to investigate the role of 11 β -HSD1 activity in macrophages. We have

used models of inflammation associated with a strong angiogenic response – the K/BxN serum transfer model of inflammatory arthritis and the sponge implantation assay, in which the role of 11 β -HSD1 activity in macrophages is examined.

Materials and methods

Animals

All experiments on animals were carried out in accordance with the UK Home Office Animals (Scientific Procedures) Act of 1986 and European Directive 2010/63/EU, following approval by the University of Edinburgh Animal Welfare and Ethical Review Body. Mice were housed in groups (2–5 per cage) under controlled conditions: 12h light/darkness cycle at 21°C with free access to standard rodent chow and water. *Hsd11b1*^{ff} mice, with *LoxP* sites flanking exon 3 of the *Hsd11b1* gene, were generated by TaconicArtemis (Cologne, Germany) onto a C57BL/6 background (Verma M, Kipari TMJ, Zhang Z, Man TY, Forster T, Homer NZM, Seckl JR, Holmes MC & Chapman KE, unpublished observations). *Hsd11b1*^{MDKO} mice with myeloid cell 11 β -HSD1 deficiency were generated by crossing *LysM-Cre* (backcrossed to a C57BL/6 background (Cramer *et al.* 2003)) with *Hsd11b1*^{ff} mice. Experimental (*Hsd11b1*^{MDKO}) and littermate control (*Hsd11b1*^{ff}) mice were the offspring of male *Hsd11b1*^{MDKO} mice bred with female *Hsd11b1*^{ff} mice. *Hsd11b1*^{Del1/Del1} mice (Verma M, Kipari TMJ, Zhang Z, Man TY, Forster T, Homer NZM, Seckl JR, Holmes MC & Chapman KE, unpublished observations) were generated from *Hsd11b1*^{ff} mice and are homozygous for a germline deletion of exon 3 of the *Hsd11b1* gene. All experiments used male mice aged between 8 and 16 weeks.

Peritoneal myeloid cell isolation and 11 β -HSD1 activity assay

Mice were killed by CO₂ asphyxiation. Resident cells were harvested from the peritoneum by lavage with 5 mL ice-cold PBS as previously described (Gilmour *et al.* 2006). Inflammatory cells elicited to the peritoneum 24 h, 72 h or 96 h following intra-peritoneal injection of thioglycollate (0.2 mL, 10%) were collected by lavage with 5 mL ice-cold PBS (Coutinho *et al.* 2016). At 24 h, cells comprise mainly neutrophils and monocytes, whereas the population is predominantly macrophages at 72 h and 96 h (Melnicoff *et al.* 1989, Zhang *et al.* 2008,

Coutinho *et al.* 2016). Where appropriate, neutrophils were isolated with anti-Ly6G antibody (clone RB6-8C5, Thermo Fisher Scientific) coupled to anti-rat IgG microbeads (Miltenyi Biotec Ltd., Bisley, Surrey, UK) by magnetic-activated cell sorting as described previously (Coutinho *et al.* 2016). Macrophages were purified by adherence to tissue culture plates using an established procedure that gives >90% purity (Zhang *et al.* 2008). 11 β -HSD1 activity in cultured cells was measured by addition of 5 nM [³H]-11-dehydrocorticosterone to the medium, as described previously (Coutinho *et al.* 2016).

K/BxN serum transfer-induced arthritis

K/BxN serum was generated in-house as described previously (Coutinho *et al.* 2012). To induce arthritis in mice, 100 μ L serum was injected intraperitoneally. Inflammation was scored daily for 21 days by an investigator blind to genotype. Scoring was carried out by visual examination according to a clinical index in which each joint was ascribed a score of 0–3, as described previously (Coutinho *et al.* 2012). The combined scores of all 4 limbs were calculated for each mouse. For histology, joints were fixed in 10% formalin for 1 day and then decalcified in 10% EDTA in neutral buffered formalin and paraffin embedded. Joint sections (4 μ m) were deparaffinised, hydrated and stained with haematoxylin and eosin for histopathological examination. The severity of pathological changes in the ankle joint was quantified on a scale of 0–3 for each mouse, based on a scoring system modified from a previous protocol (Ruiz-Heiland *et al.* 2012). Briefly, the main pathological feature, synovium thickening, is assessed and assigned a score of 0 if there is no change compared to untreated, 1 if there is mild proliferation, 2 if there is extended proliferation and 3 if there is severe proliferation with tenosynovitis. RNA was extracted from the hind ankle joints using Trizol (Thermo Fisher Scientific) following pulverisation using a mortar and pestle, under liquid nitrogen.

Sponge implant assay of angiogenesis

Mice were anaesthetised with isoflurane, and a sterilized sponge cube measuring 1 \times 1 \times 1 cm (Caligen Foam, Accrington, Lancashire, UK) was implanted subcutaneously on each flank, as described previously (Small *et al.* 2005). After 21 days, mice were killed and sponges were removed. The sponge from the right flank was frozen at 80°C for later RNA extraction using Trizol (Thermo Fisher Scientific). The sponge from the left flank was fixed in 10% formalin and embedded

in paraffin. Sections were stained with haematoxylin and eosin for angiogenesis scoring. Vessel density was determined by Chalkley counting, as described previously (Small *et al.* 2005) by an investigator blind to experimental group. Briefly, a 25-point Chalkley eyepiece graticule (Graticules Ltd, Edenbridge, Kent, UK) was used to count blood vessels at \times 250 magnification. The graticule was placed so that the maximum number of graticule dots overlay with blood vessels. The average count values (3 random areas/section, 2 sections/sample) were recorded as vessel numbers.

Immunohistochemistry

Paraffin-embedded sections (4 μ m) were deparaffinised, blocked with 10% goat serum after antigen retrieval with citrate buffer (pH 6) and incubated overnight with primary antibodies against isolectin IB4 (Alexa Fluor 488-conjugated isolectin B4, 1:100; I21411, Thermo Fisher Scientific) and α -smooth muscle actin (anti-actin, α -smooth muscle-Cy3 antibody, mouse monoclonal clone 1A4; C6198-100UL, Sigma-Aldrich) to detect endothelial cells and perivascular mural cells, respectively.

Western blotting

Peritoneal cells or purified Ly6G⁺ cells were homogenised in RIPA buffer (R0278, Sigma-Aldrich) supplemented with proteinase inhibitor cocktail (1:100 dilution; P8340, Sigma-Aldrich) and phosphatase inhibitor (1:100 dilution; P2850, Sigma-Aldrich). Proteins (20 μ g/sample) were separated by electrophoresis on a 4–12% Bis-Tris gel (NP0323BOX, Thermo Fisher Scientific) and then transferred to a 0.4 μ m nitrocellulose membrane. Antibodies used for Western blotting recognise 11 β -HSD1 (raised in sheep and kindly provided by Dr Scott Webster, the University of Edinburgh (De Sousa Peixoto *et al.* 2008, Coutinho *et al.* 2016)), β -tubulin (MAB3408, Merck Millipore) and GAPDH (ab9485, Abcam).

RNA analysis

RNA (1 μ g) was reverse transcribed into cDNA using a SuperScript III Reverse transcriptase system kit (Thermo Fisher Scientific). Levels of specific cDNAs were measured by quantitative (q)PCR in triplicate, using a LightCycler 480 (Roche) and Universal Probe Library (UPL; Roche)-based assays. Primer and probe information is shown in [Supplementary Table 1](#) (see section on [supplementary data](#) given at the end of this article). A standard curve

was prepared from pooled cDNA samples. Relative quantification was provided by LightCycler software using the maximum second derivative method and mRNA levels were normalised to an internal standard (chosen according to invariance between groups and validated against 18S RNA); *Tbp* or *Hprt*, as indicated in figure legends.

Statistics

Values are means \pm standard error of means (S.E.M.). Data were analysed using GraphPad Prism 5.0. Unpaired *t*-tests (with Welch's corrections when variance was unequal), and one-way and two-way analysis of variance (ANOVA) were used, as appropriate. Significance was set at $P < 0.05$.

Results

Disruption of 11 β -HSD1 in resident macrophages

To generate *Hsd11b1*^{MKO} mice with disruption of 11 β -HSD1 expression in myeloid phagocytes, *LysM-Cre* transgenic mice (Cramer *et al.* 2003) were crossed with mice in which exon 3 of the *Hsd11b1* gene is flanked by *LoxP* sites (*Hsd11b1*^{ff} mice) (recombination strategy shown in Supplementary Fig. 1). *LysM-Cre* transgenic mice are reported to efficiently delete *LoxP*-flanked target genes in granulocytes and mature macrophages, with lower efficiencies in other myeloid cells (Clausen *et al.* 1999).

Measurement of 11 β -HSD1 activity in myeloid cell populations from *Hsd11b1*^{MKO} mice showed an 89% decrease in 11 β -HSD1 activity in peritoneal resident cells, compared to *Hsd11b1*^{ff} littermate controls (Fig. 1A), consistent with the disruption of *Hsd11b1* in this predominantly macrophage population (Clausen *et al.* 1999). In contrast, there was no significant reduction in 11 β -HSD1 activity in thioglycollate-elicited peritoneal cells from *Hsd11b1*^{MKO} mice collected 24 h (monocytes/macrophages and neutrophils (Melnicoff *et al.* 1989, Coutinho *et al.* 2016)) or 96 h following thioglycollate injection (predominantly macrophages (Melnicoff *et al.* 1989, Zhang *et al.* 2008)) (Fig. 1B and C). Similarly, 11 β -HSD1 activity was not significantly altered in purified neutrophils (Ly6G⁺ cells) from the peritoneum of thioglycollate-injected *Hsd11b1*^{MKO} mice, compared to littermate controls (Fig. 1D). 11 β -HSD1 activity was maintained in these myeloid cell populations despite the efficient reduction in *Hsd11b1* mRNA levels (Supplementary Fig. 2). Western blotting confirmed a marked reduction in 11 β -HSD1 protein levels in the resident peritoneal cell population of *Hsd11b1*^{MKO} mice, but only a modest reduction in peritoneal cells collected 24 h or 96 h following thioglycollate injection and in purified neutrophils (Supplementary Fig. 3). These data suggest that the protein half-life of 11 β -HSD1 exceeds the half-life of the thioglycollate-elicited monocyte/macrophage and neutrophil populations, but that 11 β -HSD1 activity is markedly reduced in the longer-lived resident




Figure 1

11 β -HSD1 activity is markedly reduced in resident peritoneal macrophages from *Hsd11b1*^{MKO} mice. Resident peritoneal cells, or cells elicited to the peritoneum by i.p. injection of 0.2 mL 10% thioglycollate (TG), were harvested from *Hsd11b1*^{MKO} (MKO: white bars) and control *Hsd11b1*^{ff} mice (Con: black bars). 11 β -HSD1 activity in peritoneal cells was measured by conversion of [³H]-11-dehydrocorticosterone to corticosterone in resident peritoneal cells (A) and in cells elicited to the peritoneum 24 h (B) or 96 h (C) following thioglycollate injection. Activity was also measured in purified neutrophils (Ly6G⁺ cells) isolated from the peritoneum 24 h after thioglycollate injection (D). 11 β -HSD1 activity is expressed as pmol corticosterone/h/10⁶ cells. Values are means \pm S.E.M. and were analysed by unpaired *t*-test ($n = 3-8$, * $P < 0.05$).

macrophage population. Given this, we sought to test the role of macrophage 11 β -HSD1 in models of inflammation in which the role of resident macrophages may be evident.

Resolution of experimental arthritis is impaired in *Hsd11b1*^{MKO} mice

Tissue-resident macrophages have been implicated in the development and resolution of inflammatory arthritis in mice induced by injection of K/BxN serum (Misharin *et al.* 2014). We therefore investigated whether macrophage 11 β -HSD1 deficiency contributes to the more severe arthritis that develops in mice globally deficient in 11 β -HSD1 following injection of arthritogenic K/BxN serum (Coutinho *et al.* 2012). *Hsd11b1*^{MKO} mice did not replicate the earlier onset of arthritis seen previously with global 11 β -HSD1 deficiency (Coutinho *et al.* 2012). However, resolution of arthritis was impaired in *Hsd11b1*^{MKO} mice, compared to *Hsd11b1*^{fl/fl} controls (Fig. 2A and B). The course of the resolution phase in *Hsd11b1*^{MKO} mice was very similar to that of *Hsd11b1*^{Del1/Del1} mice globally deficient in 11 β -HSD1 (Fig. 2A and B). Histopathological examination of the joints showed

bone erosion and marked fibroproliferation in the synovium (Fig. 2C). Quantification of histopathology using a scoring index showed more marked histopathological changes in *Hsd11b1*^{MKO} mice, compared to littermate controls (Fig. 2D).

Rheumatoid arthritis is associated with neo-angiogenesis within the affected joint (Jackson *et al.* 1997, Koch 1998). Consistent with this, immunofluorescent staining of joints for the endothelial cell marker, isolectin IB4, and α -smooth muscle actin (α -SMA), a marker of perivascular mural cells, revealed a high degree of vascularisation of the soft tissues of joints of both *Hsd11b1*^{MKO} and control mice following K/BxN serum transfer arthritis (Fig. 3A). However, qPCR measurement of *Cdh5* mRNA encoding the endothelial marker, vascular endothelial (VE)-cadherin, showed >2-fold higher levels of *Cdh5* mRNA in joints of *Hsd11b1*^{MKO} mice, compared to littermate controls (Fig. 3B).

Greater inflammatory angiogenesis in *Hsd11b1*^{MKO} mice following subcutaneous sponge implantation

Increased levels of *Cdh5* mRNA in joints of *Hsd11b1*^{MKO} mice suggest an increase in neovascularisation of the joint





Figure 2


Resolution of K/BxN serum transfer-induced arthritis is impaired in *Hsd11b1*^{MKO} mice. Arthritis was induced in *Hsd11b1*^{Del1/Del1} (global knockout, GKO: triangular symbols, *n*=3), *Hsd11b1*^{MKO} (MKO: circular symbols/white bars, *n*=6) and control *Hsd11b1*^{fl/fl} mice (Con: square symbols/black bars, *n*=7) by i.p. injection of 100 μ L K/BxN serum at day 0. (A) Clinical scoring of joint inflammation over 21 days. Values are means \pm S.E.M. for *Hsd11b1*^{MKO} and control *Hsd11b1*^{fl/fl} mice. Data were analysed by two-way repeated measurement analysis of variance (ANOVA) with Bonferroni's multiple comparisons test; **P*<0.05, ***P*<0.01, ****P*<0.001. Only the mean value is shown for *Hsd11b1*^{Del1/Del1} mice. (B) The area under the curve (AUC) was calculated for the clinical score from days 9 to 21 (the resolution phase). Values are means \pm S.E.M. Data were analysed by one-way ANOVA (*P*<0.01) with *post-hoc* Dunnett's multiple comparisons test with the *Hsd11b1*^{fl/fl} group as control; **P*<0.05. (C) Representative sections of joints from *Hsd11b1*^{MKO} and *Hsd11b1*^{fl/fl} mice collected 21 days after injection and stained with haematoxylin and eosin showing tenosynovitis characterised by synovium hyperplasia, bone erosion and new bone formation. Scale bars, 100 μ m. (D) Histopathological changes were quantified using a scoring index (see the 'Materials and methods' section for details) by an investigator blind to genotype. Values are means \pm S.E.M. Data were analysed by Mann-Whitney test, *n*=6, **P*<0.05.

**Figure 3**

Higher *Cdh5* expression suggests greater angiogenesis in the joints of *Hsd11b1*^{MKO} mice. Arthritis was induced by i.p. injection of 100 μ L K/BxN serum, and the mice were killed 21 days later. (A) Immunofluorescent staining of blood vessels in the mesenchymal tissue of *Hsd11b1*^{MKO} (MKO: top) and control mice (Con: bottom). From left to right: DAPI, isolectin B4, α -SMA. (B) RNA was extracted from hind joints and qPCR used to measure levels of *Cdh5* mRNA relative to *Hprt*, used as an internal standard. Values are in arbitrary units (AU) and are means \pm S.E.M. Data from *Hsd11b1*^{MKO} (white bar) and *Hsd11b1*^{fl/fl} mice (black bar) were analysed by unpaired *t*-test, *n* = 5–7, **P* < 0.05.

during inflammatory arthritis. Mice with global 11 β -HSD1-deficiency show greater inflammatory angiogenesis induced by subcutaneous sponge implantation (Small *et al.* 2005). To test whether macrophage 11 β -HSD1 influences

inflammatory angiogenesis, sponges were implanted subcutaneously in *Hsd11b1*^{MKO} and littermate control mice, with angiogenesis assessed 21 days later (Figure 4A). Quantification of blood vessel density in sponges showed

**Figure 4**

Compared to littermate controls, *Hsd11b1*^{MKO} mice show greater angiogenesis in the subcutaneous sponge implantation assay. (A) Representative images ($\times 125$ magnification) of haematoxylin and eosin-stained sections of implanted sponges removed after 21 days, showing neovascularisation (blood vessels: arrows). (B) Quantification of blood vessel numbers by Chalkley counting. RNA was extracted from sponges removed 21 days after implantation, and qPCR was used to measure levels of: (C) *Cdh5* mRNA, (D) *Il1b* mRNA, and (E) *Angiopoietin-1*, *-2* and *-4* mRNAs, relative to levels of *Hprt* mRNA, used as an internal standard (mRNA values in arbitrary units, AU). Values are means \pm S.E.M. Data from *Hsd11b1*^{MKO} (MKO: white bars) and *Hsd11b1*^{fl/fl} mice (Con: black bars) were analysed by unpaired *t*-test (A, B, C, E) or unpaired *t*-test with Welch's correction (D); **P* < 0.05, *n* = 10–11.

20% greater vessel density in *Hsd11b1*^{MKO} mice (Fig. 4A and B) associated with almost 2-fold higher levels of *Cdh5* mRNA (Fig. 4C). Levels of mRNA encoding IL-1 β , a pro-angiogenic cytokine (Voronov *et al.* 2003), were >3-fold increased in sponges from *Hsd11b1*^{MKO} mice (Fig. 4D) and there was also a trend for increased expression of mRNA encoding pro-angiogenic angiopoietins (Fig. 4E). Consistent with previous data from mice globally deficient in 11 β -HSD1 (McSweeney *et al.* 2010), increased angiogenesis in *Hsd11b1*^{MKO} mice was not associated with a change in *Vegfa* mRNA levels, nor were levels of mRNA encoding several other cytokines altered (Supplementary Fig. 4). These data support a role for macrophage 11 β -HSD1 in controlling angiogenesis and thus promoting the resolution of inflammatory arthritis.

Discussion

Our data in *Hsd11b1*^{MKO} mice demonstrate that macrophages are a key cell type responsible for the pro-angiogenic phenotype conferred by 11 β -HSD1 deficiency. This suggests that 11 β -HSD1-mediated glucocorticoid regeneration within macrophages alters their phenotype to control an ongoing inflammatory response. In *Hsd11b1*^{MKO} mice, this happens without the perturbation in the systemic ratio of active to inactive glucocorticoid levels that occurs with global 11 β -HSD1 deficiency (Harris *et al.* 2001; Verma M, Kipari TMJ, Zhang Z, Man TY, Forster T, Homer NZM, Seckl JR, Holmes MC & Chapman KE, unpublished observations). Whether this is mediated by alterations in macrophage polarisation is currently unclear. However, *in vitro*, bone marrow-derived macrophages from *Hsd11b1*^{-/-} mice behave similar to wild type (Gilmour *et al.* 2006, Zhang & Daynes 2007), suggesting the *in vivo* environment (and possibly substrate availability) is crucial to the physiological outcomes with 11 β -HSD1 deficiency. This is suggested by evidence of an earlier macrophage polarisation to a pro-angiogenic 'M2' phenotype after myocardial infarction (McSweeney *et al.* 2010) and by higher expression of the M2 marker, SHIP1, in macrophages of *Hsd11b1*^{-/-} mice than wild-type controls (Zhang & Daynes 2007).

Whilst resident peritoneal macrophages from *Hsd11b1*^{MKO} mice showed the predicted reduction in 11 β -HSD1 activity and *Hsd11b1* mRNA, our finding of near-normal 11 β -HSD1 protein and activity in myeloid cells recruited to the peritoneum of thioglycollate-injected *Hsd11b1*^{MKO} mice, despite a marked reduction in the encoding mRNA, was unexpected. We have previously

reported a discrepancy between 11 β -HSD1 protein/activity and levels of the encoding *Hsd11b1* mRNA in mouse neutrophils, with 11 β -HSD1 protein present despite little encoding mRNA (Coutinho *et al.* 2016). Other studies have noted similar discrepancies in other cell types (Bujalska *et al.* 2005, Chinetti-Gbaguidi *et al.* 2012). This suggests that in most myeloid phagocytes 11 β -HSD1 protein has a long half-life and persists without renewal from ongoing mRNA translation, at least when inflammatory conditions prevail. The reduction in 11 β -HSD1 activity in resident peritoneal macrophages but not in thioglycollate-elicited macrophages may reflect the longer lifespan of the resident macrophage population compared to recruited bone marrow-derived macrophages (Davies *et al.* 2013). Alternatively, the *LysM-Cre* transgene may be more highly expressed in resident macrophages. Expression of *LysM/Lyz2*, encoding LYSM, is known to be heterogeneous in macrophage populations (Faust *et al.* 2000, Hume 2011). In peritoneal cells, expression of LYSM is reportedly highest in large peritoneal macrophages that are highly phagocytic (Lichanska *et al.* 1999) and correspond to the resident population (Cain *et al.* 2013). This implicates resident macrophages in the pro-angiogenic phenotype seen with 11 β -HSD1 deficiency. Resident macrophages are important in tissue repair. They efficiently clear apoptotic cells and actively promote revascularisation (Uderhardt *et al.* 2012, Wang & Kubes 2016). Tissue-resident macrophages promote vessel anastomosis and thus vascular network complexity, acting downstream of vascular endothelial growth factor (VEGF)- α (Fantin *et al.* 2010). Our finding of unchanged expression of *Vegfa* in *Hsd11b1*^{MKO} compared to control mice (also unchanged in global 11 β -HSD1-deficient mice in a model of myocardial infarction (McSweeney *et al.* 2010)) is consistent with 11 β -HSD1 acting in tissue-resident macrophages to restrain angiogenesis, downstream of VEGF- α . However, VEGF signalling is regulated at the post-translational level and further investigation is required to establish whether (or how) VEGF signalling is involved in the increased angiogenesis that occurs in *Hsd11b1*^{MKO} mice. Our data also suggest resident macrophages are an important target of glucocorticoid action in the resolution of inflammation. However, it remains possible that, in addition to macrophages, other cell types contribute to the pro-angiogenic phenotype observed with global 11 β -HSD1 deficiency.

Thus, *Hsd11b1*^{MKO} mice support a role for 11 β -HSD1 in restraining and shaping the healing response orchestrated by resident macrophages in ischaemic or injured tissue. These mice will be a useful tool with which to dissect the contribution of macrophages to beneficial (or otherwise)

effects of 11 β -HSD1 in other models of tissue repair and/or chronic inflammation.

Supplementary data

This is linked to the online version of the paper at <http://dx.doi.org/10.1530/JOE-17-0223>.

Declaration of interest

J R S holds patents on selective 11 β -HSD1 inhibitors. None of the other authors has a conflict of interest.

Funding

This work was supported by an MRC project grant (number 86642) and also by a Wellcome Trust programme grant (number 083184/Z/07/Z). Z Z was supported by a China Scholarships Council/University of Edinburgh Scholarship.

Acknowledgements

We are grateful to staff at the University of Edinburgh Central Bioresearch Services for assistance with animal care. We thank colleagues in the Centres for Cardiovascular Science and Inflammation Research for helpful advice and discussion.

References

- Bhattacharyya S, Brown DE, Brewer JA, Vogt SK & Muglia LJ 2007 Macrophage glucocorticoid receptors regulate Toll-like receptor-4-mediated inflammatory responses by selective inhibition of p38 MAP kinase. *Blood* **109** 4313–4319. (doi:10.1182/blood-2006-10-048215)
- Bujalska IJ, Draper N, Michailidou Z, Tomlinson JW, White PC, Chapman KE, Walker EA & Stewart PM 2005 Hexose-6-phosphate dehydrogenase confers oxo-reductase activity upon 11 β -hydroxysteroid dehydrogenase type 1. *Journal of Molecular Endocrinology* **34** 675–684. (doi:10.1677/jme.1.01718)
- Cain DW, O'Koren EG, Kan MJ, Womble M, Sempowski GD, Hopper K, Gunn MD & Kelsoe G 2013 Identification of a tissue-specific, C/EBP β -dependent pathway of differentiation for murine peritoneal macrophages. *Journal of Immunology* **191** 4665–4675. (doi:10.4049/jimmunol.1300581)
- Chapman KE, Coutinho AE, Zhang Z, Kipari T, Savill JS & Seckl JR 2013a Changing glucocorticoid action: 11 β -hydroxysteroid dehydrogenase type 1 in acute and chronic inflammation. *Journal of Steroid Biochemistry and Molecular Biology* **137** 82–92. (doi:10.1016/j.jsbmb.2013.02.002)
- Chapman KE, Holmes MC & Seckl JR 2013b 11 β -Hydroxysteroid dehydrogenases: intracellular gate-keepers of tissue glucocorticoid action. *Physiological Reviews* **93** 1139–1206. (doi:10.1152/physrev.00020.2012)
- Chinetti-Gbaguidi G, Bouhrel MA, Copin C, Duhem C, Derudas B, Neve B, Noel B, Eckhout J, Lefebvre P, Seckl JR, *et al.* 2012 Peroxisome proliferator activated receptor- γ activation induces 11 β -hydroxysteroid dehydrogenase type 1 activity in human alternative macrophages. *Arteriosclerosis, Thrombosis, and Vascular Biology* **32** 677–685. (doi:10.1161/ATVBAHA.111.241364)
- Clausen BE, Burkhardt C, Reith W, Renkawitz R & Forster I 1999 Conditional gene targeting in macrophages and granulocytes using LysMcre mice. *Transgenic Research* **8** 265–277. (doi:10.1023/A:1008942828960)
- Coutinho AE & Chapman KE 2011 The anti-inflammatory and immunosuppressive effects of glucocorticoids, recent developments and mechanistic insights. *Molecular and Cellular Endocrinology* **335** 2–13. (doi:10.1016/j.mce.2010.04.005)
- Coutinho AE, Gray M, Brownstein DG, Salter DM, Sawatzky DA, Clay S, Gilmour JS, Seckl JR, Savill JS & Chapman KE 2012 11 β -Hydroxysteroid dehydrogenase type 1, but not type 2, deficiency worsens acute inflammation and experimental arthritis in mice. *Endocrinology* **153** 234–240. (doi:10.1210/en.2011-1398)
- Coutinho AE, Kipari TM, Zhang Z, Esteves CL, Lucas CD, Gilmour JS, Webster SP, Walker BR, Hughes J, Savill JS, *et al.* 2016 11 β -Hydroxysteroid dehydrogenase type 1 is expressed in neutrophils and restrains an inflammatory response in male mice. *Endocrinology* **157** 2928–2936. (doi:10.1210/en.2016-1118)
- Cramer T, Yamanishi Y, Clausen BE, Forster I, Pawlinski R, Mackman N, Haase VH, Jaenisch R, Corr M, Nizet V, *et al.* 2003 HIF-1 α is essential for myeloid cell-mediated inflammation. *Cell* **112** 645–657. (doi:10.1016/S0092-8674(03)00154-5)
- Davies LC, Jenkins SJ, Allen JE & Taylor PR 2013 Tissue-resident macrophages. *Nature Immunology* **14** 986–995. (doi:10.1038/ni.2705)
- De Sousa Peixoto RA, Turban S, Battle JH, Chapman KE, Seckl JR & Morton NM 2008 Preadipocyte 11 β -hydroxysteroid dehydrogenase type 1 is a keto-reductase and contributes to diet-induced visceral obesity in vivo. *Endocrinology* **149** 1861–1868. (doi:10.1210/en.2007-1028)
- Fantin A, Vieira JM, Gestri G, Denti L, Schwarz Q, Prykhodzhij S, Peri F, Wilson SW & Ruhrberg C 2010 Tissue macrophages act as cellular chaperones for vascular anastomosis downstream of VEGF-mediated endothelial tip cell induction. *Blood* **116** 829–840. (doi:10.1182/blood-2009-12-257832)
- Faust N, Varas F, Kelly LM, Heck S & Graf T 2000 Insertion of enhanced green fluorescent protein into the lysozyme gene creates mice with green fluorescent granulocytes and macrophages. *Blood* **96** 719–726.
- Garcia RA, Search DJ, Lupisella JA, Ostrowski J, Guan B, Chen J, Yang WP, Truong A, He A, Zhang R, *et al.* 2013 11 β -Hydroxysteroid dehydrogenase type 1 gene knockout attenuates atherosclerosis and in vivo foam cell formation in hyperlipidemic apoE(-/-) mice. *PLoS ONE* **8** e53192. (doi:10.1371/journal.pone.0053192)
- Gilmour JS, Coutinho AE, Caillhier JF, Man TY, Clay M, Thomas G, Harris HJ, Mullins JJ, Seckl JR, Savill JS, *et al.* 2006 Local amplification of glucocorticoids by 11 β -hydroxysteroid dehydrogenase type 1 promotes macrophage phagocytosis of apoptotic leukocytes. *Journal of Immunology* **176** 7605–7611. (doi:10.4049/jimmunol.176.12.7605)
- Harris HJ, Kotelevtsev Y, Mullins JJ, Seckl JR & Holmes MC 2001 Intracellular regeneration of glucocorticoids by 11 β -hydroxysteroid dehydrogenase (11 β -HSD)-1 plays a key role in regulation of the hypothalamic-pituitary-adrenal axis: analysis of 11 β -HSD-1 deficient mice. *Endocrinology* **142** 114–120. (doi:10.1210/endo.142.1.7887)
- Hume DA 2011 Applications of myeloid-specific promoters in transgenic mice support in vivo imaging and functional genomics but do not support the concept of distinct macrophage and dendritic cell lineages or roles in immunity. *Journal of Leukocyte Biology* **89** 525–538. (doi:10.1189/jlb.0810472)
- Jackson JR, Seed MP, Kircher CH, Willoughby DA & Winkler JD 1997 The codependence of angiogenesis and chronic inflammation. *FASEB Journal* **11** 457–465.
- Kipari T, Hadoke PW, Iqbal J, Man TY, Miller E, Coutinho AE, Zhang Z, Sullivan KM, Mitic T, Livingstone DE, *et al.* 2013 11 β -Hydroxysteroid dehydrogenase type 1 deficiency in bone marrow-derived cells reduces atherosclerosis. *FASEB Journal* **27** 1519–1531. (doi:10.1096/fj.12-219105)
- Koch AE 1998 Review: angiogenesis: implications for rheumatoid arthritis. *Arthritis and Rheumatology* **41** 951–962. (doi:10.1002/1529-0131(199806)41:6<951::AID-ART2>3.0.CO;2-D)

- Lichanska AM, Browne CM, Henkel GW, Murphy KM, Ostrowski MC, McKercher SR, Maki RA & Hume DA 1999 Differentiation of the mononuclear phagocyte system during mouse embryogenesis: the role of transcription factor PU.1. *Blood* **94** 127–138.
- Luo MJ, Thieringer R, Springer MS, Wright SD, Hermanowski-Vosatka A, Plump A, Balkovec JM, Cheng K, Ding GJ, Kawka DW, *et al.* 2013 11 β -HSD1 inhibition reduces atherosclerosis in mice by altering pro-inflammatory gene expression in the vasculature. *Physiological Genomics* **45** 47–57. (doi:10.1152/physiolgenomics.00109.2012)
- McSweeney SJ, Hadoke PW, Kozak AM, Small GR, Khaled H, Walker BR & Gray GA 2010 Improved heart function follows enhanced inflammatory cell recruitment and angiogenesis in 11 β -HSD1-deficient mice post-MI. *Cardiovascular Research* **88** 159–167. (doi:10.1093/cvr/cvq149)
- Melnicoff MJ, Horan PK & Morahan PS 1989 Kinetics of changes in peritoneal cell populations following acute inflammation. *Cellular Immunology* **118** 178–191. (doi:10.1016/0008-8749(89)90367-5)
- Michailidou Z, Turban S, Miller E, Zou X, Schrader J, Ratcliffe PJ, Hadoke PW, Walker BR, Iredale JP, Morton NM, *et al.* 2012 Increased angiogenesis protects against adipose hypoxia and fibrosis in metabolic disease-resistant 11 β -hydroxysteroid dehydrogenase type 1 (HSD1)-deficient Mice. *Journal of Biological Chemistry* **287** 4188–4197. (doi:10.1074/jbc.M111.259325)
- Minutti CM, Knipper JA, Allen JE & Zaiss DM 2016 Tissue-specific contribution of macrophages to wound healing. *Seminars in Cell and Developmental Biology* **61** 3–11. (doi:10.1016/j.semcdb.2016.08.006)
- Misharin AV, Cuda CM, Saber R, Turner JD, Gierut AK, Haines GK 3rd, Berdnikovs S, Filer A, Clark AR, Buckley CD, *et al.* 2014 Nonclassical Ly6C(-) monocytes drive the development of inflammatory arthritis in mice. *Cell Reports* **9** 591–604. (doi:10.1016/j.celrep.2014.09.032)
- Nucera S, Bizziato D & De Palma M 2011 The interplay between macrophages and angiogenesis in development, tissue injury and regeneration. *International Journal of Developmental Biology* **55** 495–503. (doi:10.1387/ijdb.103227sn)
- Ruiz-Heiland G, Horn A, Zerr P, Hofstetter W, Baum W, Stock M, Distler JH, Nimmerjahn F, Schett G & Zwerina J 2012 Blockade of the hedgehog pathway inhibits osteophyte formation in arthritis. *Annals of the Rheumatic Diseases* **71** 400–407. (doi:10.1136/ard.2010.148262)
- Small GR, Hadoke PW, Sharif I, Dover AR, Armour D, Kenyon CJ, Gray GA & Walker BR 2005 Preventing local regeneration of glucocorticoids by 11 β -hydroxysteroid dehydrogenase type 1 enhances angiogenesis. *PNAS* **102** 12165–12170. (doi:10.1073/pnas.0500641102)
- Tuckermann JP, Kleiman A, Moriggl R, Spanbroek R, Neumann A, Illing A, Clausen BE, Stride B, Forster I, Habenicht AJ, *et al.* 2007 Macrophages and neutrophils are the targets for immune suppression by glucocorticoids in contact allergy. *Journal of Clinical Investigation* **117** 1381–1390. (doi:10.1172/JCI28034)
- Uderhardt S, Herrmann M, Oskolkova OV, Aschermann S, Bicker W, Ipseiz N, Sarter K, Frey B, Rothe T, Voll R, *et al.* 2012 12/15-lipoxygenase orchestrates the clearance of apoptotic cells and maintains immunologic tolerance. *Immunity* **36** 834–846. (doi:10.1016/j.immuni.2012.03.010)
- Voronov E, Shouval DS, Krelin Y, Cagnano E, Benharroch D, Iwakura Y, Dinarello CA & Apte RN 2003 IL-1 is required for tumor invasiveness and angiogenesis. *PNAS* **100** 2645–2650. (doi:10.1073/pnas.0437939100)
- Wang J & Kubes P 2016 A reservoir of mature cavity macrophages that can rapidly invade visceral organs to affect tissue repair. *Cell* **165** 668–678. (doi:10.1016/j.cell.2016.03.009)
- Zhang TY & Daynes RA 2007 Macrophages from 11 β -hydroxysteroid dehydrogenase type 1-deficient mice exhibit an increased sensitivity to lipopolysaccharide stimulation due to TGF- β -mediated up-regulation of SHIP1 expression. *Journal of Immunology* **179** 6325–6335. (doi:10.4049/jimmunol.179.9.6325)
- Zhang X, Goncalves R & Mosser DM 2008 The isolation and characterization of murine macrophages. *Current Protocols in Immunology* **Chapter 14** Unit 14 11. (doi:10.1002/0471142735.im1401s83)

Received in final form 26 June 2017

Accepted 4 July 2017

Accepted Preprint published online 4 July 2017